Synthesis some 4-substituted 9,10-anthraquinones

Vasyl I. Shupeniuk^{a*}, Tetyana M. Taras^a, Oksana P. Sabadakh^a, Eugene R. Luchkevich^a,
Yurii Eu. Kornii^b

^a*Department of Chemistry of the Environment and Chemical Education, Vasyl Stefanyk Precarpathian National University, Shevchenko Street 57, Ivano-Frankivsk 76018, Ukraine*

^b*Department of Chemistry of Bioactive Nitrogen-Containing Heterocyclic Bases, V.P. Kukhar Institute of Bioorganic Chemistry and Petrochemistry, NAS of Ukraine, Murmanska st., 1, Kyiv, 02094, Ukraine*

vasyl.shupeniuk@pnu.edu.ua

Keywords: Bromaminic acid, Ullmann reaction, LC-MS, 4-substituted 9,10-anthraquinones, diethanolamine.

New 4-substituted 9,10-anthraquinones (6 compounds) with amino derivations fragments were synthesized through the substitution of the bromaminic acid by amines using the Ullmann coupling reaction. The structures of the synthesized compounds were determined using LC-MS, ¹H NMR, ¹³C NMR spectroscopy, and elemental analysis data.

Introduction

The classical and most widely used strategy for the synthesis of 4-substituted anthraquinone derivatives include the Ullmann coupling reaction [1-4], which involves the treatment of bromaminic acid **1**, with an amine in the presence of a copper catalyst (**Scheme 1**) [5-

8]. The reaction typically requires harsh conditions, e.g., high temperatures and long reaction times [9-12]; it suffers from mostly poor yields product **2** [13-15], and the formation of side-products, such as **3** and **4** [16-17].

Scheme 1. Classical synthesis of 4-substituted anthraquinone derivatives

Glanzel M. described [18] the anisotropy effect of the aromatic anthraquinone ring C, 3-H-proton is protonation by carbon atoms C-9 or C-10. He also described the effect of the methyl groups rotation of the substituted ring-D, and the fragmentation of the 3-H proton from the structure of the molecule, as shown in **Figure 1**.

Figure 1. Structures of some 4-substituted 9,10-anthraquinone derivatives

Levsen K. [19], described the dissociation of $[\text{M}+\text{H}]^+$ ions of hydroxyethylamino-anthraquinone **5** (**Figure 2**), to the C-N bond as a result in the elimination of the hydroxyethyl radical ($^*\text{CH}_2\text{CH}_2\text{OH}$).

Figure 2. Dissociation 4-substituted 9,10-anthraquinone derivative **5**.

Mass spectrometry is a well recognized highly sensitive characterization method for 4-substituted anthraquinone derivatives [20], which we used for our study.

Experimental part

Material and methods

All materials purchased from commercial sources and used without purification. Melting points measured in open capillary tubes and uncorrected. The elemental analyses (C, H, N) performed using the Perkin-Elmer 2400 CHN analyzer and were within 0.4% of the theoretical values.

^1H , ^{13}C NMR spectra were recorded at Varian 400 spectrometer operating at 400 MHz frequency for ^1H and 100 MHz for ^{13}C experiments. The peaks were internally referenced to TMS (0.00 ppm) or to the residual undeuterated solvent signal. Peak multiplicities are reported as follows: s = singlet, br s = broad singlet, d = doublet, t = triplet, m = multiplet.

The chromatomass spectra were recorded on an Agilent 1100 Series high performance liquid chromatography equipped with a diode matrix with an Agilent LC/MS mass selective detector allowing a fast switching the positive/negative ionization modes. The reaction progress was monitored by the TLC method on Silica gel plates (DC-Fertigfolien ALUGRAM Xtra SILG/UV254, Germany).

Synthesis

General procedure for preparation of 4-substituted-9,10-anthraquinone 14-15, 17-20.

Bromaminic acid (0.01 mol), was dissolved in 40 ml hot water (70-80 °C), amino derivatives (0.01 mol), acid binding agent sodium bicarbonate (0.02 mol), copper sulfate (0.05 g) and ferrous sulfate (0.05 g) catalysts were then added to it. The reaction mixture stirred and heated to 90 °C. Maintained temperature 90 °C for 4 h under stirring. The product salted out by adding sodium chloride, cooled to room temperature, filtered and washed with 10% w/v brine solution.

1-Amino-4-[bis(2-hydroxyethyl)amino]-9,10-dioxo-9,10-dihydroanthracene-2-sulfonic acid 14. Blue solid; yield: 46%, m.p. 295-297 °C. LC/MS spectrum, m/z: 405,9 [M+H]⁺; C₁₈H₁₈N₂O₇S; Calculated m/z: 407. ¹H NMR (400 MHz, DMSO-d₆): δ 1.67 (d, 4H, NCH₂), 3.67 (d, J = 4.8 Hz, 4H, OCH₂), 7.49 (br s, 1H, H-3), 7.78 (m, 4H, ArH), 10.14 (s, 1H, OH), 10.79 (br s, 1H, OH); ¹H NMR (400 MHz, DMSO-d₆ + CCl₄): δ 1.68 (d, 4H, NCH₂), 3.48 (d, 2H, OCH₂), 3.69 (d, 2H, OCH₂), 7.38 (br s, 1H, H-3), 7.74 (d, J = 7.6 Hz, 2H, ArH), 7.85 (m, 2H, ArH), 9.35 (s, 1H, OH), 10.14 (br s, 1H, OH). ¹³C NMR (100 MHz, DMSO-d₆): 39.31, 39.5, 39.7 (CH₂); 112.8, 119.51, 120.49, 123.55, 126.22, 129.37, 132.31, 133.19, 133.81, 134.68, 136.37, 137.35 (arom.); 182.00, 184.06 (C=O). Anal. Calcd for C₁₈H₁₈N₂O₇S: C, 53.07; H 4.42; N 6.87; S 7.86. Found: C, 53.10; H 4.4; N 6.9; S 7.8.

1-Amino-4-[(2-hydroxyethyl)amino]-9,10-dioxo-9,10-dihydroanthracene-2-sulfonic acid 15. ¹H NMR (400 MHz, DMSO-d₆): δ 3.49 (d, 2H, CH₂), 3.69 (d, 2H, CH₂), 7.73 (s, 1H, H-3), 7.85 (t, J = 7.7 Hz, 2H, H-6, H-7), 8.25 (d, J = 8 Hz, 2H, H-5, H-8). ¹³C NMR (100 MHz, DMSO-d₆): 45.29 (CH₂); 60.24 (CH₂-OH); 109.38, 109.67, 121.16, 126.19, 126.36, 132.88, 133.03, 134.43, 134.47, 143.5, 143.79, 145.84 (arom.); 181.17, 182.12 (C=O). Anal. Calcd for C₁₈H₁₈N₂O₇S: C, 50.78; H 4.0; N 7.29; S 8.04. Found: C, 50.40; H 3.87; N 7.4; S 8.24.

4-[(4-Amino-9,10-dioxo-3-sulfo-9,10-dihydroanthracen-1-yl)-amino]butanoic acid 17. Blue solid; yield 56%, m.p. 301-303 °C. LC/MS spectrum: Found, m/z: 405,0 [M+H]⁺; C₁₈H₁₆N₂O₇S; Calculated m/z: 405. ¹H NMR (400 MHz, DMSO-d₆): δ 1.76 (d, 2H, NCH₂), 1.91 (d, 2H, CH₂), 2.32 (m, 2H, CH₂), 2.79 (s, 2H, OCH₂), 7.04 (s, 1H, H-3), 7.78 (s, 1H, ArH), 8.04 (s, 1H, ArH), 8.13 (s, 1H, ArH), 8.22 (s, 1H, ArH); ¹H NMR (400 MHz, DMSO-d₆ + CCl₄): δ 1.82 (d, 2H, NCH₂), 1.98 (d, 2H, CH₂), 2.32 (d, 2H, CH₂), 2.83 (d, 2H, CH₂), 7.2 (br s, 1H, H-3), 7.73 (d, 2H, ArH), 8.05 (d, J = 8 Hz, 2H, ArH). ¹³C NMR (100 MHz, DMSO-d₆): 40.14, 42.24, 49.47, 53.22 (CH₂); 108.23, 113.48, 125.18, 125.9, 133.05, 134.73, 140.98, 143.98, 144.34, 145.34, 148.43, 154.95 (arom.); 181.43, 186.69 (C=O). Anal. Calcd for C₁₈H₁₆N₂O₇S: C, 53.33; H 3.95; N 6.91; S 7.9. Found: C, 53.3; H 3.9; N 6.9; S 8.1.

1-Amino-4-(1H-imidazol-1-yl)-9,10-dioxo-9,10-dihydroanthracene-2-sulfonic acid **18**. Yellow solid; Recr. from aceton yield 100%, m.p. 255-257 °C. LC/MS spectrum: Found, m/z : 370,2 $[M+H]^+$; $C_{17}H_{11}N_3O_5S$; Calculated m/z : 370. 1H NMR (400 MHz, DMSO- d_6): δ 7.8 (m, $J = 8$ Hz, 8H, ArH), 9.53 (s, 1H, SO_3H). Anal. Calcd for $C_{17}H_{11}N_3O_5S$: C, 55.13; H 2.97; N 11.35; S 8.65. Found: C, 55.10; H 2.9; N 11.37; S 8.63.

1-Amino-9,10-dioxo-4-(2-oxo-4-sulfanylidene-1,3-thiazolidin-3-yl)-9,10-dihydroanthracene-2-sulfonic acid **19**. Blue solid; Yield 14.3%, m.p. 285-288 °C. LC/MS spectrum: Found, m/z : 435,2 $[M+H]^+$; $C_{17}H_{11}N_3O_5S$; Calculated m/z : 434. 1H NMR (400 MHz, DMSO- d_6): δ 3.32 (d, 2H, CH_2), 7.26 (s, 1H, H-3), 7.82 (d, $J = 7.2$ Hz, 2H, H-6, H-7), 8.33 (d, $J = 8$ Hz, 2H, H-5, H-8). Anal. Calcd for $C_{17}H_{10}N_2O_6S_3$: C, 46.89; S 22.06. Found: C, 46.85; S 22.00.

1-Amino-4-[(pyridin-3-yl)amino]-9,10-dioxo-9,10-dihydroanthracene-2-sulfonic acid **20**. Blue solid; Yield 15%, m.p. 250-251 °C. LC/MS spectrum: Found, m/z : 394,2 $[M+H]^+$; $C_{19}H_{13}N_3O_5S$; Calculated m/z : 395. 1H NMR (400 MHz, DMSO- d_6): δ 7.62 (s, 1H, H-3), 7.7 (d, 2H, H-5, H-7), 7.85 (t, $J = 6.4$ Hz, 3H, ArH), 8.08 (d, $J = 6.4$ Hz, 2H, H-5, H-8); 1H NMR (400 MHz, DMSO- d_6 + CCl_4): δ 7.63 (br s, 1H, H-3), 7.84 (t, $J = 6.8$ Hz, 2H, H-6, H-7), 7.96 (s, 1H, ArH), 8.00 (s, 1H, ArH), 8.10 (s, 1H, ArH), 8.17 (d, $J = 6.8$ Hz, 2H, H-5, H-8). ^{13}C NMR (100 MHz, DMSO- d_6): 122.58, 124.42, 127.11,

127.40, 131.71, 133.22, 134.83, 135.58, 136.29, 140.89, 152.98, 173.58 (arom.); 180.82, 181.09 (C=O). Anal. Calcd for $C_{19}H_{13}N_3O_5S$: C, 57.73; H 3.29; N 10.63; S 8.1. Found: C, 57.7; H 3.3; N 10.7; S 8.17.

Results and discussion

Substitution with dialkyl amines **6-7** was starting according to the procedure described previously [21]. However, the desirable products were not obtained. In the case of nucleophilic substitution on diethanolamine (7) formed product **15** - 37% on the LC-MS (m/z 363.2 $[M+H]^+$). In the case of substitution brom on diethylamine (6), formations of side-product (4) was 28% (m/z 302.0 $[M+H]^+$). Then we used the general procedure (experimental data) for nucleophilic substitution in bromaminic acid (**Scheme 3**). In the case of substitution brom of diethanolamine (7), the product **15** was still present in LC-MS (m/z 364.0 $[M+H]^+$). The ^{13}C NMR spectra characteristic signals carbon (CH_2-CH_2-OH) was 45.29 and 60.24 ppm; the resonance signals for two carbonyl carbons 1-amino-4-[(2-hydroxyethyl)amino]-9,10-dioxo-9,10-dihydro-anthracene-2-sulfonic acid (**15**) at δ 181.17 and 182.12 ppm. Ullmann coupling reaction of bromaminic acid with diethylamine (6), leads to the formation product **13** (m/z 347.0 $[M+H]^+$), it was fixed using diode array detection (DAD).

The desired 4-diethanol-substituted product **14**, separated by HPLC. 1H NMR is

present in 8 aliphatic protons of methylene groups in solvents (DMSO- d_6) and (DMSO- d_6 +CCl $_4$), characteristic shifts of 5 aromatic protons at 7.38-8.24 ppm, respectively.

Figure 3. Proposed mechanism

The proposed mechanism for the formation of compound **15** (m/z 364.0 [M+H]⁺) (Figure 2), depicted in the LC-MS spectrum (Scheme 2). And explains fragmentation 3-H-proton of the ethylhydroxyl radical which transformed into oxirane.

Scheme 2. Chromatomass-spectra

substitution by amino acids

In case substitution brom of glycine **8**, is formation of side-products **4** - 38.9% (m/z 318.2 [M+H]⁺) and **3** was 24.5% (m/z 302.0 [M+H]⁺). The poor yields of product **16** explained by the weak base of aminoacetic acid (**8**). Reaction of bromaminic acid with amino acid **9**, the formation of isomers **17** (m/z 405.0 [M+H]⁺).

After purification by silica gel flash column chromatography, eluted with ethyl acetate:acetone 1:3 v/v, compound **17** was obtained. The ¹H NMR spectrum in (DMSO- d_6) there are 6 aliphatic CH₂ methylene protons in the region at 1.76-2.79 ppm. The ¹³C NMR spectra was characteristic signals carbon methylene groups in the region at 40.14-53.22 ppm.

substitution by heterocyclic amines

The yield of product **18** after recrystallization with acetone was 100% (m/z 370.2 [M+H]⁺). The aromatic protons were visible as a multiplet of δ 7.7-7.8 ppm. In the case of substitution bromaminic acid with 3-aminopyridine (**12**), the yield of the main product **20** was 14.9% (m/z 394.0 [M+H]⁺) the ¹H NMR spectrum nine aromatic protons appeared in the region of 7.62-8.24 ppm (DMSO- d_6) and 7.63-8.17 (DMSO- d_6 + CCl $_4$).

Scheme 3. Synthesis of 4-substituted 9,10-anthraquinonesulfonic acids

The main undesirable product identified in all reactions was 1-amino-4-hydroxy-9,10-dioxo-9,10-dihydroanthracene 2-sulfonate (**4**, **Scheme 3**) formed by attack of the competing nucleophile hydroxide.

Conclusions

Fragmentation mechanisms of protonated 4-substituted-9,10-anthraquinones and its derivatives investigated by atmospheric pressure ionization with mass spectrometry (MS). The major fragmentation pathways were loss of the

hydroxyethyl group (*CH_2CH_2OH) or ethyl group (*CH_2CH_3) from fourth position of the **C** ring. Determined LC/MS that as a result of nucleophilic substitution of brom on γ -aminobutyric acid, the final products are the isomers of compound **17**.

Acknowledgements

The work was carried out with the financial support of the Ministry of Education and Science of Ukraine (Project No. 0119U103131).

References

- [1] Ullmann F.; Bielecki J. Ueber Synthesen in der Biphenylreihe". *Chemische Berichte.*, 1901, 34(2): 2174–2185. doi:10.1002/cber.190103402141
- [2] Jiao, J.; Zhang, X.-R.; Chang, N.-H.; et.al. A facile and practical copper powder-catalyzed, organic solvent-and ligand-free Ullmann amination of aryl halides. *J. Org. Chem.* 2011, 76:1180-1183; DOI: 10.1021/jo102169t
- [3] Ma, D.W.; Cai, Q.; Copper/amino acid catalyzed cross-couplings of aryl and vinyl halides with nucleophiles *Acc. Chem. Res.*, 2008, 41:1450-1460; DOI: 10.1021/ar8000298
- [4] Carril, M.; Martin, R.S.; Dominguez, E.; Palladium and copper-catalysed arylation reactions in the presence of water, with a focus on carbon-heteroatom bond formation. *Chem. Soc. Rev.*, 2008, 37:639 - 647; DOI: 10.1039/B709565C
- [5] Patel, N.B.; Patel, A.L.; Characterization, application and microbial study of imidazole base acid anthraquinone dyes. *Oriental J. of Chem.*, 2008, 24:551-558.
- [6] Glanzel, M.; Bultmann, R.; Starke, K.; Frahm, A.W. Constitutional isomer of Reactive Blue 2 – selective P2Y-receptor antagonists ? *Eur. J. Med. Chem.* 2003, 38:303–312.
- [7] Glanzel, M.; Bultmann, R.; Starke, K.; Frahm, A.W. Structure-activity relationships of novel P2-receptor antagonists structurally related to Reactive Blue 2. *Eur. J. Med. Chem.* 2005, 40:1262–1276.
- [8] Baqi Y.; Muller C.E. Synthesis of alkyl- and aryl-amino-substituted anthraquinone derivatives by microwave-assisted copper (0)-catalyzed Ullmann coupling reactions. *Nature Prot.* 2010, 5:387-390; DOI: 10.1038/nprot.2010.63
- [9] Lee, C.-C.; Chen, C.-L.; Liu, F.-L.; et. al. Development of 1-Amino-4-(phenylamino)anthraquinone-2-sulfonate Sodium Derivatives as a New Class of Inhibitors of RANKL-Induced Osteoclastogenesis. *Arch. Pharm. Chem. Life Sci.*, 2016, 349:342–355. DOI: 10.1002/ardp.201500475
- [10] Baqi Y.; Weyler S.; Iqbal J.; Muller C.E. Structure-activity relationships of anthraquinone derivatives derived from bromaminic acid as inhibitors of ectonucleoside triphosphate diphosphohydrolases (E-NTPDases). *Purinergic Signalling.* 2009, 5:91–106; DOI: 10.1007/s11302-008-9103-5
- [11] Baqi Y.; Malik E.; Rashed M.; Wingen L.; Muller C.E. Ullmann reactions of 1-amino-4-bromoanthraquinone bearing various 2-substituents furnishing novel dyes. *Dyes and Pigments.* 2016, 131:33-40; DOI: 10.1016/j.dyepig.2016.03.023
- [12] Baqi Y.; Malik E.; Muller C.E. Syntheses of 2-substituted 1-amino-4-bromoanthraquinones (bromaminic acid analogues) – precursors for dyes and drugs. *Beilstein J. Org. Chem.*, 2015, 11:2326–2333; DOI: 10.3762/bjoc.11.253
- [13] Roy S.; Large R.J.; Akande A.M.; et.al. Development of GoSlo-SR-5-69, a potent activator of large conductance Ca²⁺-activated K⁺ (BK) channels. *Eur. J. Med. Chem.*, 2014, 75:426–437; DOI: org/10.1016/j.ejmech.2014.01.035
- [14] Olubunmi, A.; Olatunji A. Gabriel, Condensation and Cellulosic Derivatization of Bromamine Acid. *The Pacific Journal of Science and Technology.* 2009, 10:788–794.
- [15] Yildiz, E.; Keles, M.; Kaya, A.; Dincer, S. Mononuclear Fe (III), Cr (III), Co (II) Metal Complexes Based on Azo-Anthraquinone Moieties: Synthesis, Characterizations and Antibacterial Activities. *Chem. Sci. Trans.*, 2013, 2(2):547–555.
- [16] Baqi Y.; Muller C.E. Rapid and efficient microwave-assisted copper (0)-catalyzed Ullmann coupling reaction. *Org. Lett.* 2007, 9:1271–1274; DOI: 10.1021/ol070102v
- [17] Vrba, Z.; Mechanism of the Ullmann condensation. *Chem. Commun*, 1981, 46:92 – 100.
- [18] Glanzel M. Synthesis P2-Pezceptor-Antagonistenin der Reiheder Reaktivblau 2 Farbstobbe

und deren Struktur-Wirkung Beziehungen: dissertation zur Erlangung der Doktorwürde. 2003.

[19] Levsen K.; Schiebel H.; Terlouw J. Even-electron ions: a systematic study of the neutral species lost in the dissociation of quasi-molecular ions. *J. Mass Spektrom.* 2007, 42:1028 – 1033. DOI: 10.1002/jms.1234

[20] Enaud, E.; Trovaslet, M.; Bruvneel, F.; *et. al.*, A novel azoanthraquinone dye made through innovative enzymatic process. *Dyes and Pigments.*, 2010, 85(3):102. DOI 10.1016/j.dyepig.2009.10.010

[21] Inoue, H.; Nakamura, K.; Kato, S.; Hida, M. The Nucleophilic Photo-substitution reaction of anthraquinone derivatives. II. The Mechanism of the Photoamination of sodium 1-amino-4-bromanthraquinone-2-sulfate. *J. Chem. Soc., of Japan.*, 1975, 48(10):2872 - 2876.